

RESEARCH

Open Access



Overexpression of IκBα in cardiomyocytes alleviates hydrogen peroxide-induced apoptosis and autophagy by inhibiting NF-κB activation

Min Han^{1,2†}, Xiao-Cui Chen^{1†}, Ming-Hui Sun³, Min-Tao Gai¹, Yi-Ning Yang², Xiao-Ming Gao¹, Xiang Ma², Bang-Dang Chen^{1*} and Yi-Tong Ma^{1,2*}

Abstract

Background: Inflammation and oxidative stress play predominant roles in the initiation and progression of ischaemia/reperfusion (I/R) injury, with nuclear factor kappa B (NF-κB) serving as a crucial mediator. Overexpression of the inhibitor of κB alpha (IκBα) gene is hypothesized to have protective effects against apoptosis and autophagy in cardiomyocytes subjected to hydrogen peroxide (H₂O₂) by inhibiting the NF-κB pathway.

Methods: The IκBα^{S32A, S36A} gene was transfected via adeno-associated virus serotype 9 (AAV9) delivery into neonatal rat ventricular cardiomyocytes (NRVMs) prior to H₂O₂ treatment. NRVMs were divided into control, H₂O₂, GFP + H₂O₂, IκBα+H₂O₂, and pyrrolidine dithiocarbamate (PDTC) + H₂O₂ groups. Nuclear translocation of the NF-κB p65 subunit was evaluated by immunofluorescence and Western blotting. Cell viability was assessed by Cell Counting Kit-8 assay. Supernatant lactate dehydrogenase (LDH) and intracellular malondialdehyde (MDA) were measured to identify H₂O₂-stimulated cytotoxicity. Apoptosis was determined by Annexin V-PE/7-AAD staining, and the mitochondrial membrane potential (ΔΨm) was detected by JC-1 staining. Western blotting was used to detect apoptosis- and autophagy-related proteins.

Results: IκBα transfection significantly increased cell viability and ΔΨm but decreased the supernatant LDH and cellular MDA levels in cardiomyocytes exposed to H₂O₂. Meanwhile, IκBα overexpression decreased H₂O₂-induced apoptosis by upregulating the Bcl-2/Bax ratio and reduced autophagy by downregulating the expression of Beclin-1 and the LC3-II/LC3-I ratio. These effects partly accounted for the ability of IκBα to inhibit the NF-κB signalling pathway, as evidenced by decreases in p65 phosphorylation and nuclear translocation. Indeed, the effects of inactivation of NF-κB signalling with the specific inhibitor PDTC resembled the cardioprotective effects of IκBα during H₂O₂ stimulation.

(Continued on next page)

* Correspondence: chenbangdang@126.com; myt_xj@sina.com

†Min Han and Xiao-Cui Chen are contributed equally to this article and share first authorship.

¹Xinjiang Key Laboratory of Cardiovascular Disease Research, Clinical Medical Research Institute, the First Affiliated Hospital of Xinjiang Medical University, Urumqi 830054, PR China

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

Conclusion: I κ B α overexpression can ameliorate H₂O₂-induced apoptosis, autophagy, oxidative injury, and $\Delta\Psi_m$ loss through inhibition of the NF- κ B signalling pathway. These findings suggest that I κ B α transfection can result in successful resistance to oxidative stress-induced damage by inhibiting NF- κ B activation, which may provide a potential therapeutic target for the prevention of myocardial I/R injury.

Keywords: Nuclear factor kappa B, Inhibitor of kappa B alpha, Apoptosis, Autophagy, Adeno-associated virus serotype 9, Oxidative stress, Cardiomyocytes

Introduction

Acute myocardial infarction (AMI) is the leading cause of death worldwide, and reperfusion therapy is the most effective treatment for AMI [1]. Paradoxically, the process of myocardial reperfusion also induces a series of adverse cardiac events such as inflammation, necrosis, apoptosis and autophagy, ultimately leading to myocardial ischaemia/reperfusion (I/R) injury [2]. Recent evidence has suggested that excessive inflammation and oxidative stress play predominant roles in the initiation and progression of I/R injury [3, 4].

Nuclear factor kappa B (NF- κ B) is an inflammatory inducer and redox-sensitive transcription factor in most cell types [5]. The p65/50 heterodimer, the most common pattern of NF- κ B dimer, normally exists as a component of inactive cytoplasmic complexes bound to the inhibitor of κ B alpha (I κ B α). Upon stimulation, I κ B α is phosphorylated and undergoes ubiquitylation and proteasomal degradation, subsequently leading to phosphorylation and nuclear translocation of the NF- κ B p65 subunit [6]. Activated NF- κ B then initiates the expression of corresponding target genes, many of which may regulate apoptosis, inflammation and autophagy [7].

However, whether NF- κ B is protective or detrimental for cardiomyocyte apoptosis remains controversial [8]. Notably, our previous study indicated that the p65 ribozyme could prevent cell apoptosis in H9C2 cardiomyocytes exposed to hydrogen peroxide (H₂O₂) [9]. Autophagy, an evolutionarily conserved form of “self-digestion”, plays dual roles in the heart [10]. Recent studies on autophagy have shown both the protective [11] and deleterious [12] effects of autophagy in cardiomyocytes against oxidative stress. Evidence has revealed a strong correlation between modulation of NF- κ B and the autophagic response [13, 14]. In addition, cross-talk between autophagy and apoptosis has been noted [15], and NF- κ B is known to mediate the balance between autophagy and apoptosis [16].

Therefore, NF- κ B activation is thought to be the key point of I/R injury; thus, inhibiting NF- κ B may be a targeted therapy for I/R injury. Phosphorylation of I κ B α , the key inhibitor of the canonical NF- κ B pathway, at Ser 32 and Ser 36 is necessary for its degradation, and any mutation of these two serine residues blocks I κ B α

degradation [6]. Recently, adeno-associated virus serotype 9 (AAV9) was demonstrated to be the best gene carrier due to its high efficiency in the heart [17]. H₂O₂, a common reactive oxygen species (ROS), is generally utilized to mimic I/R injury in vitro [12]. Thus, the I κ B α ^{S32A, S36A} gene was transfected into cardiomyocytes via AAV9-mediated delivery to investigate the role of inhibition of the NF- κ B pathway in H₂O₂-induced apoptosis and autophagy. Pyrrolidine dithiocarbamate (PDTC), a specific inhibitor of NF- κ B, was used as a positive control in this study.

Materials and methods

Ethics statement

The experimental protocol was approved by the Ethics Committee of the First Affiliated Hospital of Xinjiang Medical University (No. IACUC-20180223-69). One- to three-day-old neonatal Sprague-Dawley (SD) rats were purchased from the Experimental Animal Center of Xinjiang Medical University and handled in accordance with the recommendations in the Guidelines for the Care and Use of Laboratory Animals of the National Institutes of Health.

Reagents

Briefly, rabbit anti-Bax polyclonal antibody (#2772) and rabbit anti-p65 (#8242), anti-p-p65 (#3033), anti-I κ B α (#4812), anti-GFP (#2956), anti-Bcl-1 (#3495), and anti-LC3 II/I (#12741) monoclonal antibodies were all obtained from Cell Signalling Technology (Danvers, MA, USA). Rabbit anti-Bcl-2 (ab196495), anti-Histone H3 (ab1791) and anti- β -actin (ab8227) polyclonal antibodies and horseradish peroxidase (HRP)-conjugated anti-rabbit secondary antibody (ab205718) were obtained from Abcam (Cambridge, UK). RIPA buffer and Halt™ Protease and Phosphatase Inhibitor Cocktail were obtained from Thermo Fisher Scientific (Waltham, MA, USA). Enhanced chemiluminescence (ECL) reagent and JC-1 were obtained from Millipore (Bedford, MA, USA). Trypsin, PDTC and bromodeoxyuridine (BrdU) were obtained from Sigma (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM), foetal bovine serum (FBS), and penicillin-streptomycin solution were obtained from Gibco (Grand Island, NY, USA). Collagenase

II was obtained from Worthington (Minnesota, USA). H_2O_2 was obtained from Sangon (Shanghai, China).

Vector design

Recombinant AAV-9 vectors generated by a recombinant baculovirus (rBac)-based system in SF9 cells as previously described were purchased from Virovek (Hayward, CA, USA) [18]. The recombinant AAV9 vectors were packaged as double-stranded DNA and contained the enhanced green fluorescent protein (eGFP) gene (dsAAV9-GFP) or the $I\kappa B\alpha^{S32A, S36A}$ gene (dsAAV9- $I\kappa B\alpha$) driven by the human cytomegalovirus (CMV) promoter.

Isolation and culture of rat cardiomyocytes

The protocol for the isolation and purification of neonatal rat ventricular cardiomyocytes (NRVMs) was reported in our previous study [19]. Briefly, the hearts of 1- to 3-day-old neonatal SD rats were dissected and digested with 0.1% trypsin and 0.08% collagenase II. Following differential adhesion twice for 50 min each time, nonadherent cells were resuspended and cultivated in high-glucose DMEM containing 10% FBS, 1% penicillin-streptomycin, and 0.1 mM BrdU for 48 h. The medium was replaced every 48 h.

AAV9 transfection of cardiomyocytes

After 48 h of culture, NRVMs were transfected with dsAAV9-GFP or dsAAV9- $I\kappa B\alpha$ as previously described [12]. Briefly, cells were first transfected with dsAAV9 (multiplicity of infection, $MOI = 5 \times 10^6$ vg/cell) in serum-free medium, and then DMEM at an equal volume containing 20% FBS, 2% penicillin-streptomycin and 0.2 mM BrdU was added to every dish 3 h later. Images showing GFP were captured using a fluorescence inverted microscope (Leica DMI4000B, Wetzlar, Germany), and the green fluorescence intensities were analysed using ImageJ software (National Institutes of Health, NY, USA).

Experimental design and cell grouping

The experiment was designed to explore whether AAV9-delivered $I\kappa B\alpha^{S32A, S36A}$ gene transfection could protect cardiomyocytes against H_2O_2 -induced apoptosis and autophagy via inhibition of NF- κB activation. Cardiomyocytes were starved with serum-free DMEM for 12 h to ensure cell synchronization before H_2O_2 stimulation. The experimental cardiomyocytes were randomly divided into 5 groups as follows: (1) the control group, which contained primary cardiomyocytes cultivated under normal conditions; (2) the H_2O_2 control group (H_2O_2): the model control group, which contained primary cardiomyocytes subjected to 100 μM H_2O_2 alone [12]; (3) the GFP control group (GFP): the vector control group, which contained primary cardiomyocytes

transfected with dsAAV9-GFP virus for 5 days before being subjected to 100 μM H_2O_2 ; (4) the $I\kappa B\alpha$ treatment group ($I\kappa B\alpha$): the treatment group, which contained primary cardiomyocytes transfected with dsAAV9- $I\kappa B\alpha$ virus for 5 days before being subjected to 100 μM H_2O_2 ; and (5) the PDTC treatment group (PDTC): the positive control group, which contained primary cardiomyocytes pretreated with 100 μM PDTC for 60 min before being subjected to 100 μM H_2O_2 .

Measurement of cardiomyocyte viability and cytotoxicity

The Cell Counting Kit-8 (CCK-8; Dojindo, Japan) assay was used to assess cell viability. In brief, 2×10^4 cells were seeded into each well of a 96-well plate and transfected with GFP or $I\kappa B\alpha$ for 5 days. After the cells were exposed to H_2O_2 , 10 μL of CCK-8 stock solution was added to each well, followed by incubation at 37 °C for 2 h. The absorbance at 450 nm was measured with a GO microplate spectrophotometer (Thermo Fisher Scientific). The extent of cell death was determined by quantifying lactate dehydrogenase (LDH) released into the culture supernatant with an LDH kit (Jiancheng Bioengineering Institute, Nanjing, China). Intracellular malondialdehyde (MDA), an indicator of oxidative injury, was also measured using an MDA assay kit (Jiancheng Bioengineering Institute).

Flow cytometry analysis

Cell apoptosis was measured using PE Annexin V Kit I (BD Biosciences, NJ, USA). Briefly, cells were collected and resuspended in 1 \times binding buffer. Then, the solution (1×10^5 cells) was supplemented with 5 μL of PE Annexin V and 7-AAD and incubated in the dark for 15 min at room temperature. Apoptotic cells were identified by flow cytometry (Beckman Coulter, CA, USA). All the experiments were performed in triplicate.

Western blotting analysis

Nuclear and cytoplasmic proteins were extracted following the instructions of a Nuclear and Cytoplasmic Extraction Kit (Thermo Fisher Scientific, USA). Total proteins were extracted with RIPA buffer containing Halt™ Protease and Phosphatase Inhibitor Cocktail. Phosphorylated p65 in the total lysate and the nuclear p65 to cytosolic p65 ratio were both detected to identify activation of the NF- κB signalling pathway [20]. Equal amounts of protein were loaded and separated on pre-cast SDS-PAGE gels (Invitrogen, Grand Island, NY, USA) and transferred to Millipore PVDF membranes. After blocking with 5% skim milk, the membranes were blotted overnight with specific primary antibodies against p65 (1:1000), p-p65 (1:500), $I\kappa B\alpha$ (1:1000), Bax (1:1000), GFP (1:1000), Beclin-1 (1:1000), LC3 II/I (1:1000), Bcl-2 (1:1000), Histone H3 (1:1000), and β -actin

(1:1000) at 4 °C, followed by incubation with anti-rabbit HRP secondary antibody (1:5000) at room temperature for 2 h. ECL solution was added to the membranes to visualize signals. β -actin and Histone H3 were regarded as loading controls. Images were captured and analysed by Image Lab 4.0 software (Bio-Rad Laboratories, Hercules, CA, USA).

Immunofluorescence

Immunofluorescence was employed to identify H_2O_2 -induced nuclear translocation of the NF- κ B p65 subunit in cardiomyocytes. Briefly, 2×10^5 cells were seeded into confocal dishes. After H_2O_2 treatment, cardiomyocytes were fixed with 4% paraformaldehyde for 20 min and permeabilized with 0.25% Triton X-100 for 10 min. After blocking with 1% BSA for 1 h, cells were probed overnight with anti-p65 antibody (1:200) at 4 °C and incubated with Alexa Fluor 594-labelled secondary antibody (Invitrogen, 1:200, labelled with red fluorescence) for 2 h at room temperature, followed by 10 min of DAPI staining of nuclei (labelled with blue fluorescence). Signals were detected using a confocal spectral microscope (Leica SP8, Germany).

Measurement of the mitochondrial membrane potential

JC-1 is an ideal fluorescent probe used to detect the mitochondrial membrane potential ($\Delta\Psi_m$) in cardiomyocytes. Briefly, a 10 nmol/L JC-1 working solution was prepared prior to use, and cardiomyocytes were stained at 37 °C in the dark for 15 min. Cells doubly stained with JC-1 were visible by either green or red fluorescence. Fluorescence images and intensities were obtained using a fluorescence microscope and ImageJ software. Generally, $\Delta\Psi_m$ is represented by the red to green fluorescence ratio, which decreases in proportion with the severity of cell injury.

Statistical analysis

All statistical analyses were performed with SPSS 22.0 (SPSS, Inc., Chicago, IL, USA). Data are presented as the mean \pm SEM. Multiple comparisons were carried out using one-way analysis of variance (ANOVA) followed by Bonferroni's post hoc test. A value of $P < 0.05$ indicated statistical significance.

Results

H_2O_2 -induced activation of NF- κ B in NRVMs

The results indicated that H_2O_2 elicited time-dependent I κ B α degradation and p65 translocation after the NRVMs were incubated with 100 μ M H_2O_2 for different durations (0, 15, 30, and 60 min). (Fig. 1a-c). The ratio of nuclear p65 to cytosolic p65 peaked at 60 min. Consistent with the nuclear translocation of p65, the level of p-p65/p65 increased following H_2O_2 stimulation with

the incubation time (Fig. 1d and e) and was highest at 60 min. Thus, treatment with 100 μ M H_2O_2 for 60 min was used in the following experiments.

Efficiency of I κ B α transfection in NRVMs

As shown in Fig. 1f, the green fluorescence signal was robust, and the dsAAV9-GFP transfection efficiency in NRVMs reached more than 70%. Western blotting analysis showed that the GFP protein was more highly expressed in the GFP group than in the other groups, while the I κ B α protein level was significantly elevated in the I κ B α group compared with the control and GFP groups (Fig. 1g-i).

I κ B α protected cardiomyocytes from H_2O_2 -induced apoptosis


The proportion of apoptotic cells in the control group was $7.0 \pm 1.5\%$. After treatment with 100 μ M H_2O_2 , the apoptotic rates of cardiomyocytes in the H_2O_2 group and GFP group increased to $21.20 \pm 0.95\%$ and $19.97 \pm 0.97\%$, respectively, which were decreased by I κ B α or PDTTC pretreatment (Fig. 2a). Indeed, compared with the levels in the control group, the anti-apoptotic protein Bcl-2 was downregulated, but the pro-apoptotic protein Bax was upregulated in NRVMs exposed to H_2O_2 , leading to a higher Bax/Bcl-2 ratio, but this effect was completely abolished by pretreatment with I κ B α or PDTTC (Fig. 2b).

I κ B α protected cardiomyocytes from H_2O_2 -induced cell injury

Compared to that in the control group, $\Delta\Psi_m$ was significantly decreased in the H_2O_2 control group, but this decrease was rescued by I κ B α or PDTTC treatment (Fig. 3a). Additionally, H_2O_2 treatment significantly decreased cell viability but elevated supernatant LDH and intracellular MDA levels; however, these changes were reversed by I κ B α or PDTTC treatment (Fig. 3b-d).

I κ B α suppressed H_2O_2 -induced NF- κ B activation and autophagy in NRVMs

Compared with the control group, H_2O_2 treatment significantly elicited p65 translocation, and these changes were successfully reversed by I κ B α or PDTTC pretreatment (Fig. 4a and b). Consistently, H_2O_2 increased p-p65/p65 ratio in NRVMs, but I κ B α or PDTTC dramatically downregulated the H_2O_2 -induced expression of p-p65. Meanwhile, Beclin-1 and the LC3-II/LC3-I ratio, the autophagy-associated markers, were markedly upregulated in NRVMs exposed to H_2O_2 , whereas these effects were inhibited by I κ B α or PDTTC treatment (Fig. 4c).




Discussion

This study shows that $I\kappa B\alpha$ degradation and NF- κB activation occurred in a time-dependent manner in NRVMs subjected to H_2O_2 . Cells treated with H_2O_2 showed reductions in cell viability and $\Delta\Psi_m$ but elevations in LDH and MDA levels, apoptosis and autophagy. $I\kappa B\alpha$ transfection or PDTC pretreatment ameliorated H_2O_2 -induced cell injury by inhibiting NF- κB activation.

I/R injury severely attenuates the benefit of revascularization after AMI and has therefore become an important focus of cardiovascular research [2]. The inflammatory response induced by AMI is essential for heart repair, but excessive generation of ROS and inflammation following reperfusion therapy exacerbate heart damage [21].

The NF- κB signalling pathway plays key roles in the inflammatory response, oxidative stress, apoptosis, and autophagy in the heart [8]. Phosphorylation and nuclear translocation of the p65 subunit are signs of NF- κB activation [20]. Previous studies [22–24] identified that H_2O_2 treatment for different durations (30 min–24 h) elicited significant p65 phosphorylation and nuclear translocation in NRVMs. In line with these studies, p65 was time-dependently phosphorylated and translocated from the cytoplasm to the nucleus with $I\kappa B\alpha$ degradation in NRVMs subjected to H_2O_2 .

However, whether NF- κB activation protects or damages cardiomyocytes remains debatable. An early study demonstrated that activation of NF- κB reduced cell apoptosis in hypoxic cardiomyocytes [25], whereas most



recent studies [26, 27] have shown that NF-κB is a proapoptotic transcription factor correlated with myocardial injury and that blockade of NF-κB activity prevents myocardial apoptosis. Gray et al. [22] recently reported that ROS generated by ischaemia-reperfusion could rapidly activate calmodulin kinase II (CaMKII), which increased cell injury by inducing IκBα degradation and nuclear p65 accumulation in NRVMs exposed to H₂O₂. Importantly, knockout of the CaMKIIδ gene significantly attenuated myocardial infarct size by inhibiting IκBα degradation and NF-κB activation. All these findings show that NF-κB activation deteriorates the heart in I/R injury.

Herein, we hypothesized that direct overexpression of IκBα to prevent NF-κB activation may have a good protective effect in cardiomyocytes. Then, dsAAV9-IκBα^{Ser32A,36A} was designed to prevent IκBα degradation due to its phosphorylation at the Ser 32 and Ser 36 sites and successfully transfected into cardiomyocytes. Western blotting and immunofluorescence demonstrated that IκBα transfection successfully maintained cytoplasmic IκBα levels and suppressed p65 phosphorylation and translocation in NRVMs exposed to H₂O₂. Additionally, IκBα elevated cell viability, decreased LDH and MDA levels, and attenuated apoptosis, implying the protective role of IκBα in H₂O₂-induced cell injury in NRVMs. The mechanisms may account for the role of NF-κB in

mediating the expression of various proteins that promote or inhibit apoptosis. Notably, NF-κB regulates the expression of certain anti-apoptotic genes, such as Bcl-2 [28], and an increased Bcl-2/Bax ratio decreases cell apoptosis. In this study, treatment with IκBα or PDTC significantly elevated the Bcl-2/Bax ratio in NRVMs subjected to H₂O₂. These data indicate that IκBα protects NRVMs against H₂O₂-induced apoptosis by increasing the Bcl-2/Bax ratio.

Opening of the mitochondrial permeability transition pore (MPTP) in the first few minutes of reperfusion leads to ΔΨ_m loss and is responsible for necrotic and apoptotic cell death processes, contributing differentially to myocardial infarct size [29]. Importantly, inhibition of the opening of the MPTP attenuated I/R injury. Thus, ΔΨ_m loss reflects mitochondrial dysfunction, indicates early-stage apoptosis and is a critical determinant of I/R injury [30]. A previous study demonstrated that H₂O₂ induced a significant decrease in ΔΨ_m [12]. In this study, H₂O₂ treatment attenuated ΔΨ_m and enhanced Bax expression in NRVMs, and these effects were reversed by pretreatment with IκBα or PDTC. NF-κB is involved in the regulation of mitochondrial dysfunction [31], and Bax antagonizes the anti-apoptotic ability of Bcl-2 and simultaneously promotes permeability of the mitochondrial outer membrane and reduces the level of




Fig. 3 IκBα reduced cell injury in NRVMs exposed to H₂O₂. **a** Representative images of JC-1 staining in different groups (scale bar: 250 μm). The results indicated that IκBα or PDTC elevated the ratio of red to green fluorescence intensity (n = 3, * P < 0.05 and ** P < 0.01 vs. Control, # P < 0.05 vs. H₂O₂ group). **b** CCK-8 results demonstrated that IκBα or PDTC increased the cell viability of NRVMs stimulated by H₂O₂ (n = 3, * P < 0.05 and ** P < 0.01 vs. Control, # P < 0.05 vs. H₂O₂ group). **c** IκBα or PDTC reduced the levels of supernatant LDH in NRVMs subjected to H₂O₂ (n = 3, * P < 0.05 and ** P < 0.01 vs. Control, # P < 0.05 and ## P < 0.01 vs. H₂O₂ group). **d** IκBα or PDTC decreased the content of intracellular MDA in H₂O₂-treated NRVMs (n = 3, ** P < 0.01 vs. Control, # P < 0.05 vs. H₂O₂ group)

$\Delta\Psi_m$ [32]. The results herein suggest that IκBα decreases cell injury and apoptosis by inhibiting NF-κB activation and Bax expression, ultimately elevating $\Delta\Psi_m$ after H₂O₂ stimulation.

Autophagy, a cellular process of lysosome-mediated degradation of cytoplasmic


components or damaged organelles, is thought to be an adaptive response and protective for cell survival [10]. However, autophagy causes a redox effect in cardiomyocytes when exposed to different stimuli. Evidence supports the benefit of autophagy to cardiomyocytes during myocardial ischaemia through its improvement of myocardial energy metabolism and organelle recycling [33], but excessive autophagy causes lethal damage to cells during cardiac I/R injury [21], which is mediated in part by upregulation of Beclin-1 expression [34].

However, the communication between autophagy and NF-κB is bidirectional. Autophagy is required for activation of NF-κB [13]; in turn, NF-κB further increases autophagosome maturation by upregulating Beclin-1 and LC3-II expression in I/R injury [35]. Importantly, PDTC attenuates Beclin-1 expression and the formation of autophagosomes by suppressing I/R injury-induced

NF-κB activation [16]. In accordance with these findings, treatment of NRVMs with H₂O₂ induced p65 phosphorylation and translocation, enhanced Beclin-1 expression, and increased the LC3-II/LC3-I ratio, and these effects were rescued by IκBα transfection or PDTC treatment. These results imply that IκBα can protect cardiomyocytes by inhibiting H₂O₂-induced autophagy. In addition, Bcl-2 can bind Beclin-1 to inhibit autophagy [36]. This study also demonstrates that IκBα transfection elevated the expression of Bcl-2, which may disturb the function of Beclin-1 and thus further inhibit H₂O₂-induced autophagy, implying cross-talk between apoptosis and autophagy.

Study strengths and limitations

The present study has several strengths. To the best of our knowledge, this study is the first to report the cardioprotective effects of IκBα transfection through inhibition of H₂O₂-induced apoptosis and autophagy. Moreover, AAV9 vectors, which can be successfully transfected into cardiomyocytes, may be used as an effective tool for gene therapy for AMI. This study also has some limitations. The present findings were derived



from neonatal cardiomyocytes in vitro and may differ from findings in animal experiments due to the complicated features of the in vivo environment. Further animal studies should be conducted to confirm the cardioprotective effects of IκBα against I/R injury.

Conclusions

The findings of this study show that pretreatment with dsAAV9-IκBα or PDTC protected NRVMs from H₂O₂-induced apoptosis, autophagy, mitochondrial dysfunction, and oxidative damage by restraining the NF-κB signalling pathway, suggesting that IκBα transfection can

protect cardiomyocytes against cardiac oxidative damage. Thus, AAV9 vectors, as high-efficiency gene carriers to heart, may be used to carry the IκBα gene to protect the heart through targeted inhibition of myocardial NF-κB in future preclinical or clinical studies, which may provide a promising gene therapy for preventing cardiac I/R injury.

Abbreviations

NF-κB: Nuclear factor kappa B; IκBα: Inhibitor of kappa B alpha; NRVMs: Neonatal rat ventricular cardiomyocytes; I/R: Ischaemia/reperfusion; AMI: Acute myocardial infarction; H₂O₂: Hydrogen peroxide; GFP: Green fluorescent protein; PDTC: Pyrrolidine dithiocarbamate; CCK-8: Cell counting

kit-8; LDH: Lactate dehydrogenase; MDA: Malondialdehyde; MPTP: Mitochondrial permeability transition pore; $\Delta\Psi_m$: Mitochondrial membrane potential; ROS: Reactive oxygen species

Acknowledgements

The authors thank Xue Zhang and Ning Yang for their technical assistance in this work.

Authors' contributions

MH, XCC, MHS, and MTG performed the research. YNY, XMG, and XM provided guidance throughout the whole study. MH and XCC analysed the data and drafted the paper. YTM and BDC designed and provided final approval of the version to be published. The author(s) read and approved the final manuscript.

Funding

This work was supported by National Natural Science Foundation of China (No. 81873490, No. 81860072), the Xinjiang Uygur Autonomous Region Graduate Research and Innovation Project (No. XJ2019G183), the Xinjiang Medical University Graduate Innovation and Entrepreneurship Project (No. CXCY2018020), and the Tianshan Innovative Research Team Plan (No. 2018D14006).

Availability of data and materials

All data generated or analysed during this study are included in this article.

Ethics approval and consent to participate

All animal experimental protocols were approved by the Ethics Committee of the First Affiliated Hospital of Xinjiang Medical University (No. IACUC-20180223-69).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Xinjiang Key Laboratory of Cardiovascular Disease Research, Clinical Medical Research Institute, the First Affiliated Hospital of Xinjiang Medical University, Urumqi 830054, PR China. ²Department of Cardiology, the First Affiliated Hospital of Xinjiang Medical University, Urumqi 830054, PR China. ³Department of Nephrology, Fifth Affiliated Hospital of Xinjiang Medical University, Urumqi 830000, PR China.

Received: 26 February 2020 Accepted: 16 June 2020

Published online: 24 June 2020

References

- Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the task force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2018;39(2):119–77.
- Del Re DP, Amgalan D, Linkermann A, Liu Q, Kitsis RN. Fundamental mechanisms of regulated cell death and implications for heart disease. *Physiol Rev*. 2019;99(4):1765–817.
- Verma VK, Malik S, Mutneja E, Sahu AK, Bhatia J, Arya DS. Attenuation of ROS-mediated myocardial ischemia-reperfusion injury by morin via regulation of RISK/SAPK pathways. *Pharmacol Rep*. 2020. <https://doi.org/10.1007/s43440-019-00011-2>.
- Wallert M, Ziegler M, Wang X, Maluenda A, Xu X, Yap ML, et al. α -Tocopherol preserves cardiac function by reducing oxidative stress and inflammation in ischemia/reperfusion injury. *Redox Biol*. 2019;26:101292.
- Sies H, Berndt C, Jones DP. Oxidative stress. *Annu Rev Biochem*. 2017;86:715–48.
- Zhang Q, Lenardo MJ, Baltimore D. 30 years of NF- κ B: a blossoming of relevance to human pathobiology. *Cell*. 2017;168(1–2):37–57.
- Sun SC. The non-canonical NF- κ B pathway in immunity and inflammation. *Nat Rev Immunol*. 2017;17(9):545–58.
- Kumar R, Yong QC, Thomas CM. Do multiple nuclear factor kappa B activation mechanisms explain its varied effects in the heart? *Ochsner J*. 2013;13(1):157–65.
- Sun Z, Ma YT, Chen BD, Liu F. Recombinant adeno-associated virus serotype 9 with p65 ribozyme protects H9c2 cells from oxidative stress through inhibiting NF- κ B signaling pathway. *J Geriatr Cardiol*. 2014;11(4):311–5.
- Sciarretta S, Maejima Y, Zablocki D, Sadoshima J. The role of autophagy in the heart. *Annu Rev Physiol*. 2018;80:1–26.
- Zhu HH, Wang XT, Sun YH, He WK, Liang JB, Mo BH, et al. Pim1 overexpression prevents apoptosis in Cardiomyocytes after exposure to hypoxia and oxidative stress via upregulating cell autophagy. *Cell Physiol Biochem*. 2018;49(6):2138–50.
- Sun MH, Chen XC, Han M, Yang YN, Gao XM, Ma X, et al. Cardioprotective effects of constitutively active MEK1 against H₂O₂-induced apoptosis and autophagy in cardiomyocytes via the ERK1/2 signaling pathway. *Biochem Biophys Res Commun*. 2019;512(1):125–30.
- Criollo A, Chereau F, Malik SA, Niso-Santano M, Marino G, Galluzzi L, et al. Autophagy is required for the activation of NF κ B. *Cell Cycle*. 2012;11(1):194–9.
- Wang X, Tao YL, Huang YW, Zhan KG, Xue M, Wang Y, et al. Catalase ameliorates diabetes-induced cardiac injury through reduced p65/RelA-mediated transcription of BECN1. *J Cell Mol Med*. 2017;21(12):3420–34.
- Dong Y, Chen H, Gao J, Liu Y, Li J, Wang J. Molecular machinery and interplay of apoptosis and autophagy in coronary heart disease. *J Mol Cell Cardiol*. 2019;136:27–41.
- Zeng M, Wei X, Wu Z, Li W, Li B, Zhen Y, et al. NF- κ B-mediated induction of autophagy in cardiac ischemia/reperfusion injury. *Biochem Biophys Res Commun*. 2013;436(2):180–5.
- García-Olloqui P, Rodríguez-Madoz JR, Di Scala M, Abizanda G, Vales Á, Olagüe C, et al. Effect of heart ischemia and administration route on biodistribution and transfection efficiency of AAV9 vectors. *J Tissue Eng Regen Med*. 2020;14(1):123–34.
- Chen H. Exploiting the Intron-splicing Mechanism of Insect Cells to Produce Viral Vectors Harboring Toxic Genes for Suicide Gene Therapy. *Mol Ther Nucleic Acids*. 2012;1(11):e57.
- Tao J, Chen BD, Ma YT, Yang YN, Li XM, Ma X, et al. FrzA gene protects cardiomyocytes from H₂O₂-induced oxidative stress through restraining the Wnt/frizzled pathway. *Lipids Health Dis*. 2015;14:90.
- Prabhu SD, Frangogiannis NG. The biological basis for cardiac repair after myocardial infarction: from inflammation to fibrosis. *Circ Res*. 2016;119(1):91–112.
- Huang Z, Han Z, Ye B, Dai Z, Shan P, Lu Z, et al. Berberine alleviates cardiac ischemia/reperfusion injury by inhibiting excessive autophagy in cardiomyocytes. *Eur J Pharmacol*. 2015;762:1–10.
- Gray CB, Suetomi T, Xiang S, Mishra S, Blackwood EA, Glembotski CC, et al. CaMKII δ subtypes differentially regulate infarct formation following ex vivo myocardial ischemia/reperfusion through NF- κ B and TNF- α . *J Mol Cell Cardiol*. 2017;103:48–55.
- Wei C, Li L, Kim IK, Sun P, Gupta S. NF- κ B mediated miR-21 regulation in cardiomyocytes apoptosis under oxidative stress. *Free Radic Res*. 2014;48(3):282–91.
- Tang B, Ma J, Ha X, Zhang Y, Xing Y. Tumor necrosis factor- α upregulated PHLPP1 through activating nuclear factor- κ B during myocardial ischemia/reperfusion. *Life Sci*. 2018;207:355–63.
- Misra A, Haudek SB, Knuefermann P, Vallejo JG, Chen ZJ, Michael LH, et al. Nuclear factor- κ B protects the adult cardiac myocyte against ischemia-induced apoptosis in a murine model of acute myocardial infarction. *Circulation*. 2003;108(25):3075–8.
- Zou G, Zhong W, Wu F, Wang X, Liu L. Inhibition of lncRNA Neat1 by catalpol via suppressing transcriptional activity of NF- κ B attenuates cardiomyocyte apoptosis. *Cell Cycle*. 2019;18(24):3432–41.
- Zhang W, Zhang Y, Ding K, Zhang H, Zhao Q, Liu Z, et al. Involvement of JNK1/2-NF- κ Bp65 in the regulation of HMGB2 in myocardial ischemia/reperfusion-induced apoptosis in human AC16 cardiomyocytes. *Biomed Pharmacother*. 2018;106:1063–71.
- Liu Q, Liu Z, Zhou LJ, Cui YL, Xu JM. The long noncoding RNA NKILA protects against myocardial ischaemic injury by enhancing myocardin expression via suppressing the NF- κ B signalling pathway. *Exp Cell Res*. 2020;387(2):111774.
- Moriciano G, Bonora M, Campo G, Aquila G, Rizzo P, Giorgi C, et al. Mechanistic role of mPTP in ischemia-reperfusion injury. *Adv Exp Med Biol*. 2017;982:169–89.

30. Zhao Y, Guo Y, Chen Y, Liu S, Wu N, Jia D. Curculigoside attenuates myocardial ischemia-reperfusion injury by inhibiting the opening of the mitochondrial permeability transition pore. *Int J Mol Med*. 2020;45(5):1514–24.
31. Han Y, Cai Y, Lai X, Wang Z, Wei S, Tan K, et al. lncRNA RMRP Prevents Mitochondrial Dysfunction and Cardiomyocyte Apoptosis via the miR-1-5p/hsp70 Axis in LPS-Induced Sepsis Mice. *Inflammation*. 2020;43(2):605–18.
32. Garcaí-Sáez AJ. The secrets of the Bcl-2 family [J]. *Cell Death Differ*. 2012; 19(11):1733–40.
33. Huang L, Dai K, Chen M, Zhou W, Wang X, Chen J, et al. The AMPK agonist PT1 and mTOR inhibitor 3HOI-BA-01 protect Cardiomyocytes after ischemia through induction of autophagy. *J Cardiovasc Pharmacol Ther*. 2016;21(1): 70–81.
34. Zuo Z, Zuo PF, Sheng ZL, Wang X, Ding JD, Ma GS. Tetramethylpazine attenuates myocardial ischemia/reperfusion injury through modulation of autophagy. *Life Sci*. 2019;239:117016.
35. Haar L, Ren X, Liu Y, Koch SE, Goines J, Tranter M, et al. Acute consumption of a high-fat diet prior to ischemia-reperfusion results in cardioprotection through NF-κB-dependent regulation of autophagic pathways. *Am J Physiol Heart Circ Physiol*. 2014;307(12):H1705–13.
36. Pattingre S, Tassa A, Qu X, Garuti R, Liang XH, Mizushima N, et al. Bcl-2 antiapoptotic proteins inhibit Beclin 1-dependent autophagy. *Cell*. 2005; 122(6):927–39.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

